

Jorge Chahla, MD, PhD

ACL Repair with Meniscal Repair (All Inside)

Post-Operative Rehabilitation Protocol

0-2 weeks

- NWB for 2 weeks
- ROM: 0-90° with emphasis on full extension
- Patella mobilization
- SLR supine with brace locked at 0 degrees
- Quad Sets with brace locked at 0 degrees
- Ankle Pumps
- Short crank (90mm) ergometry

2-6 weeks

- Begin WBAT
- No WB with the knee past 90 degrees of flexion
- Discontinue crutches when gait is non-antalgic
- ROM: 0-125 degrees (maintain full extension)
- Active knee extension to 40 degrees
- Standard (170 mm) ergometry (ROM >115°)
- Leg press (80-0 degree arc)
- Mini squats/weight shifts
- Proprioception training
- Initiate Step Up program
- Avoid tibial rotation until 6 weeks

6-14 weeks Progressive squat program

- Initiate Step-Down program
- Leg press, lunges
- Isotonic knee extensions (90-40 degrees, close chain preferred)
- Agility exercises (sport cord)
- Versaclimber /Nordic track
- Retrograde treadmill ambulation

14-22 weeks

- Begin forward running (treadmill program when 8" stepdown status satisfactory
- Continue strengthening and flexibility program
- Advance sports-specific agility drills
- Start plyometric program

Weight Bearing Status

NWB x 2 weeks

Brace Settings

Locked straight x 3-4 weeks when ambulating or sleeping. Otherwise open brace once good quad control

ROM Restrictions

Full PROM beginning POD 1



>22 weeks

- Advance plyometric program, return to sport (MD directed)
- **May perform Functional Sports Assessment (FSA) 7 months and then subsequent test at 9 months postop for clearance to return to sport

| Exercise | Week | | | | | | | | | | | | | |
|---|---------------|---|--------|-----|-----|----|------|------------------|------------|-----|-----|------|------------|----|
| | 1 2 | 3 | | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 12 | 16 | 20 | 24 |
| Initial Exercises: | II | | | | | | | | | 1 | | 1 | | |
| ROM Goals | 0-90 ° | | 0-125° | | | | | Progress to full | | | | | | |
| Flexion/extension, heel slides | | | | | | | | | Ŭ | | | | | |
| Flexion/extension- seated | | | | | | | | | | | | | | |
| Patella/Tendon mobilization | | | | | | | | | | | | | | |
| Quad Series | | | | | | | | | | | | | | |
| Hamstring Sets | | | | | | | | | | | | | | |
| Sit and Reach for Hamstrings | | | | | | | | | | | | | | |
| Ankle Pumps | | | | | | | | | | | | | | |
| Crutch weaning | NWB | | | | | | | | | | | | | |
| Heel prop knee extension stretch | | | | | | | | | | | | | | |
| SLR (w/ brace until quad control restored) | | | | | | | | | | | | | | |
| Toe and heel raises | | | | | | | | | | | | | | |
| Weight-bearing Strength Exercises: | 1 2 | 3 | | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 12 | 16 | 20 | 24 |
| Double Knee Bends | | | | | | | | | | | | | | |
| Double Leg Bridges | | | | | | | | | | | | | | |
| Step up/Lunge Progression | | | | | IS | Os | (| 0-7(|) ° | | 7 | 0-9 | D ° | |
| Beginning cord exercises | NWB | | | | | | | | | | | | | |
| Proprioception/Balance Training | | | | | | | | | | | | | | |
| Dead Lift $(2 \rightarrow 1)$ | | | | | | | | | | | | | | |
| Squat/Leg Press (ISO →reps, 2→1 leg) | | | | 0-4 | 15° | 0- | •70° | | | >70 | gra | adua | al | |
| Sports Test Exercises | | | | | | | | | | | | | | |
| Cardiovascular Exercises: | 1 2 | 3 | | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 12 | 16 | 20 | 24 |
| Bike with both legs-no resistance | | | | | | | | | | | | | | |
| Bike with both legs- resistance | | | | | | | | | | | | | | |
| Aqua jogging | | | | | | | | | | | | | | |
| Treadmill-walking 7% incline | NWB | | | | | | | | | | | | | |
| Swimming with fins-light flutter kick | | | | | | | | | | | | | | |
| Elliptical Trainer | | | | | | | | | | | | | | |
| Rowing | | | | | | | | | | | | | | |
| Stair stepper | | | | | | | | | | | | | | |
| Agility Exercises: | 1 2 | 3 | | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 12 | 16 | 20 | 24 |
| Running Progression | | | | | | | | | | | | | | |
| Initial-Single Plane | NWB | | | | | | | | | | | | | |
| Advance-Multi Directional | | | | | | | | | | | | | | |
| Functional Sports Test | | | | | | | | | | | | | | |
| High Level Activities: | 1 2 | 3 | | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 12 | 16 | 20 | 24 |
| Golf Progression | | | | | | | | | | | | | | |
| Outdoor biking, hiking, running | NWB | | | | | | | | | | | | | |
| Return to Full Sport at 9 months post-op | | | | | | | | | | | | | | |

Quad Series:

- •Quad sets (5 sec hold) also quick contractions (NMES to supplement)
- Prone quad sets
- •SLR with no extension leg
- Progression of Quad Strengthening (2-6 weeks)
- •LAQ in safe range -no added resistance to begin
- •Once weight bearing tandem stance TKE with 5 sec hold
- Isometric wall sit
- •SLR with resistance
- •Step up progression
- Squat progression
- Leg press progression
- Lunge progression

Hamstring Series:

- •Hamstring stretching (supine/prone)
- •Hamstring set (heel dig)- contract hamstring with knee flexed pressing heel into the floor (5 sec hold)
- Hamstring curls
- Progress to resisted hamstring curl

Sport test exercises:

- •Goals: impact training and introduction of directional drills
- Double leg squat jumps
- •Box jumps
- Speed skaters
- Double leg jumps for height and distance progress to single leg as tolerated
- •Blaze pod change of direction drills

Further Clarifications and Considerations:

- •Double Knee Bends: feet shoulder width apart, bend knees to 30^o, keeping knees behind the toes.
- •Cord exercises (resistance band/cord): resisted double knee bends, resisted hamstring curls
- •May begin to implement BFR as early as 1 week into post-op period
- •Functional Sports Test: Baseline completed around 7 months post-op with subsequent test at 9 months

Return to Running Criteria

- Trace effusion, flexion within 5 degrees of contralateral side
- LSI on anterior reach Y balance test \ge 90%
- LSI on quadriceps torque output on isometric measurement ≥ 75%
- 12" SL squat tolerance
- Able to walk > 1 mile with no limping or pain
- Able to "hop" with UE assist on operative leg 5 times without pain or compensation
- •SL balance with eyes closed 30 seconds without loss of balance



Return to Agility/Change of Direction Criteria

- •Return to running criteria listed above met
- •No effusion
- •Full ROM
- •Quad LSI on isokinetic ≥85%
- Hamstring LSI on isokinetic ≥85%
- •LSI on anterior reach Y-balance \geq **95%**
- •SL hopping pain-free and effusion free

Return to Sport Criteria

- In order to test must demonstrate full AROM, no pain, no/trace effusion, 5/5 strength, negative lachman and pivot shift, LSI ≥95% hamstring curl and leg press
- •Quadricep strength ≥95% of contralateral side (< 2cm of atrophy)
- Able to perform single leg squat with correct form
 LSI SL hop ≥95%
- Y-Balance ≥95% (mean of 3 trials in anterior, posterolateral, posteromedial /100)
- Vertical jump test, single leg hop distance, and timed single leg hop over 20 -feet ≥90% of contralateral side

