

Jorge Chahla, MD, PhD

Meniscal Repair (All Inside) Post-Operative Rehabilitation Protocol

0-2 weeks	 NWB for 2 weeks with brace locked in extension AROM/PROM: 0-90° with emphasis on full extension Patella mobilization SLR supine with brace locked at 0 degrees 	Weight Bearing Status
	 Quad Sets with brace locked at 0 degrees Quad Sets with brace locked at 0 degrees Heel slides Ankle Pumps 	NWB x 2 weeks
2-4 weeks	 Begin WBAT progression: advance to PWB (25% then 50%) starting at 2 weeks continue to progress Can unlock brace at this time if maintain full extension 	Brace Settings
	 No WB with the knee past 90 degrees of flexion Progress with full ROM until full (maintain full extension) 	Locked straight x 3-4 weeks when ambulating
4-8 weeks	 Advance to full WBAT with brace unlocked Discontinue brace when quad strength adequate (typically around 4 weeks) Discontinue crutches when gait normalized and non-antalgic Wall sits to 90 degrees 	or sleeping. Otherwise open brace once good quad control
8-12 weeks	 Continue with full ROM Progress with closed chain exercises Lunges from 0-70 degrees Leg press 0-70 degrees Begin stationary bike 	ROM Restrictions
12-16 weeks	 Progress strengthening exercises Single leg strengthening Begin jogging and progress to running Sports specific exercise 	beginning POD 1

_

	Week													
Exercise	1	2	3	4	5	6	7	8	9	10	12	16	20	24
Initial Exercises:		1												1
ROM Goals		.90°	0-125°				Progress to full							
Flexion/extension, heel slides		0.00		0123			Trog							
Flexion/extension- seated														
Patella/Tendon mobilization														
Quad Series														
Hamstring Sets														
Sit and Reach for Hamstrings														
Ankle Pumps														
Crutch weaning	N	NB												
Heel prop knee extension stretch														
SLR (w/ brace until quad control restored)														
Toe and heel raises														
Balance Series														
Weight-bearing Strength Exercises:	1	2	3	4	5	6	7	8	9	10	12	16	20	24
Double Knee Bends					1				<u> </u>	<u> </u>	1	1	<u> </u>	
Double Leg Bridges														
Reverse Lunge- static hold														
Beginning cord exercises	N۱	NWB												
Dead Lift $(2 \rightarrow 1)$														
Squat/Leg Press (ISO →reps, 2→1 leg)				0-45	5°			0	-70	· (70) [,] m	ax)		
Sports Test Exercises														
Cardiovascular Exercises:	1	2	3	4	5	6	7	8	9	10	12	16	20	24
Bike with both legs-no resistance														
Bike with both legs- resistance														
Aqua jogging														
Treadmill-walking 7% incline	N۱	WB												
Swimming with fins-light flutter kick														
Elliptical Trainer														
Rowing														
Stair stepper														
Agility Exercises:	1	2	3	4	5	6	7	8	9	10	12	16	20	24
Running Progression														
Initial-Single Plane	N۱	NWB												
Advance-Multi Directional														
Functional Sports Test														
High Level Activities:	1	2	3	4	5	6	7	8	9	10	12	16	20	24
Golf Progression														
Outdoor biking, hiking, running	NWB													
Return to Full Sport at 6 months post-op														

Adapted from post-operative protocol created by Howard Head Physical Therapy, Jill Monson, PT, OCS and Jon Schoenecker, PT, OCS, CSCS

Quad Series:

- •Quad sets (5 sec hold) also quick contractions (NMES to supplement)
- •Prone quad sets
- •SLR with no extension leg
- Progression of Quad Strengthening
- •LAQ in safe range -no added resistance to begin
- •Once weight bearing tandem stance TKE with 5 sec hold
- Isometric wall sit
- •SLR with resistance
- Step up progression
- Squat progression
- Leg press progression
- Lunge progression

Hamstring Series:

- •Hamstring stretching (supine/prone)
- •Hamstring set (heel dig)- contract hamstring with knee flexed pressing heel into the floor (5 sec hold)
- •Hamstring curls
- Progress to resisted hamstring curl

Sport Test Exercises:


- •Goals: impact training and introduction of directional drills
- Double leg squat jumps
- •Box jumps
- Speed skaters
- Double leg jumps for height and distance progress to single leg as tolerated
- •Blaze pod change of direction drills

Further Clarifications and Considerations

- •Double Knee Bends: feet shoulder width apart, bend knees to 30^o, keeping knees behind the toes.
- •Cord exercises (resistance band/cord): resisted double knee bends, resisted hamstring curls
- •May begin to implement BFR as early as 1 week into post-op period
- •Functional Sports Test: can be completed at around 6 months

Return to Running Criteria

- Trace effusion, flexion within 5 degrees of contralateral side
- LSI on anterior reach Y balance test \ge 90%
- LSI on quadriceps torque output on isometric measurement ≥ 75%
- 12" SL squat tolerance
- Able to walk > 1 mile with no limping or pain
- •Able to "hop" with UE assist on operative leg 5 times without pain or compensation
- •SL balance with eyes closed 30 seconds without loss of balance



- Return to running criteria listed above met
- No effusion
- •Full ROM
- •Quad LSI on isokinetic ≥85%
- Hamstring LSI on isokinetic ≥85%
- •LSI on anterior reach Y-balance \geq **95%**
- •SL hopping pain-free and effusion free

Return to Sport Criteria

- In order to test must demonstrate full AROM, no pain, no/trace effusion, 5/5 strength, negative lachman and pivot shift, LSI ≥95% hamstring curl and leg press
- •Quadricep strength ≥95% of contralateral side (< 2cm of atrophy)
- Able to perform single leg squat with correct form
 LSI SL hop ≥95%
- •Y-Balance ≥95% (mean of 3 trials in anterior, posterolateral, posteromedial /100)
- Vertical jump test, single leg hop distance, and timed single leg hop over 20 -feet ≥90% of contralateral side

