



Meniscal Repair (All Inside)

Post-Operative Rehabilitation Protocol

0-2 weeks

- NWB for 2 weeks with brace locked in extension
- AROM/PROM: 0-90° with emphasis on full extension
- Patella mobilization
- SLR supine with brace locked at 0°
- Quad Sets with brace locked at 0°
- Heel slides
- Ankle Pumps

2-4 weeks

- Begin WBAT progression: advance to PWB (25% then 50%) starting at 2 weeks continue to progress
- Can unlock brace at this time if maintain full extension
- No WB with the knee past 90° of flexion
- Progress with full ROM until full (maintain full extension)

4-8 weeks

- Advance to full WBAT with brace unlocked
- Discontinue brace when quad strength adequate (typically around 4 weeks)
- Discontinue crutches when gait normalized and non-antalgic
- Wall sits to 90°

8-12 weeks

- Continue with full ROM
- Progress with closed chain exercises
- Lunges from 0-70°
- Leg press 0-70°
- Begin stationary bike

12-16 weeks



- Progress strengthening exercises
- Single leg strengthening
- Begin jogging and progress to running
- Sports specific exercise
- Plyo Prep Screen when initiating running and/or jumping. Lower Body Assessment when initiating return to sport progression. Schedule via QR code email or spc@rushortho.com

Weight Bearing Status

NWB x 2
weeks

Brace Settings

Locked
straight x 3-4
weeks when
ambulating
or sleeping.
Otherwise
open brace
once good
quad control

ROM Restrictions

Full PROM
beginning
POD 1



Exercise	Week																	
	1	2	3	4	5	6	7	8	9	10	12	16	20	24				
Initial Exercises:																		
ROM Goals	0-90°		0-125°			Progress to full												
Flexion/extension, heel slides																		
Flexion/extension- seated																		
Patella/Tendon mobilization																		
Quad Series																		
Hamstring Sets																		
Sit and Reach for Hamstrings																		
Ankle Pumps																		
Crutch weaning	NWB																	
Heel prop knee extension stretch																		
SLR (w/ brace until quad control restored)																		
Toe and heel raises																		
Balance Series																		
Weight-bearing Strength Exercises:	1	2	3	4	5	6	7	8	9	10	12	16	20	24				
Double Knee Bends	NWB																	
Double Leg Bridges																		
Reverse Lunge- static hold																		
Beginning cord exercises																		
Dead Lift (2 → 1)																		
Squat/Leg Press (ISO →reps, 2→1 leg)			0-45				0-70 (70 max)											
Sports Test Exercises																		
Cardiovascular Exercises:			1	2	3	4	5	6	7	8	9	10	12	16	20	24		
Bike with both legs-no resistance	NWB																	
Bike with both legs- resistance																		
Aqua jogging																		
Treadmill-walking 7% incline																		
Swimming with fins-light flutter kick																		
Elliptical Trainer																		
Rowing																		
Stair stepper																		
Agility Exercises:	1	2	3	4	5	6	7	8	9	10	12	16	20	24				
Running Progression	NWB																	
Initial-Single Plane																		
Advance-Multi Directional																		
Functional Sports Test																		
High Level Activities:	1	2	3	4	5	6	7	8	9	10	12	16	20	24				
Golf Progression	NWB																	
Outdoor biking, hiking, running																		
Return to Full Sport at 6 months post-op																		



Quad Series:

- Quad sets (5 sec hold) also quick contractions (NMES to supplement)
- Prone quad sets
- SLR with no extension leg
- Progression of Quad Strengthening
 - LAQ in safe range -no added resistance to begin
 - Once weight bearing tandem stance TKE with 5 sec hold
 - Isometric wall sit
 - SLR with resistance
 - Step up progression
 - Squat progression
 - Leg press progression
 - Lunge progression

Hamstring Series:

- Hamstring stretching (supine/prone)
- Hamstring set (heel dig)- contract hamstring with knee flexed pressing heel into the floor (5 sec hold)
- Hamstring curls
- Progress to resisted hamstring curl

Sport Test Exercises:

- Goals: impact training and introduction of directional drills
- Double leg squat jumps
- Box jumps
- Speed skaters
- Double leg jumps for height and distance progress to single leg as tolerated
- Blaze pod change of direction drills

Further Clarifications and Considerations

- Double Knee Bends: feet shoulder width apart, bend knees to 30°, keeping knees behind the toes.
- Cord exercises (resistance band/cord): resisted double knee bends, resisted hamstring curls
- May begin to implement BFR as early as 1 week into post-op period
- Functional Sports Test: can be completed at around 6 months




Return to Running Criteria

- Trace effusion, flexion within 5 degrees of contralateral side
- LSI on anterior reach Y balance test $\geq 90\%$
- LSI on quadriceps torque output on isometric measurement $\geq 75\%$
- 12" SL squat tolerance
- Able to walk > 1 mile with no limping or pain
- Able to "hop" with UE assist on operative leg 5 times without pain or compensation
- SL balance with eyes closed 30 seconds without loss of balance



Return to Agility/Change of Direction Criteria

- Return to running criteria listed above met
- No effusion
- Full ROM
- Quad LSI on isokinetic $\geq 85\%$
- Hamstring LSI on isokinetic $\geq 85\%$
- LSI on anterior reach Y-balance $\geq 95\%$
- SL hopping pain-free and effusion free



Return to Sport Criteria

- In order to test must demonstrate full AROM, no pain, no/trace effusion, 5/5 strength, negative lachman and pivot shift, LSI $\geq 95\%$ hamstring curl and leg press
- Quadricep strength $\geq 95\%$ of contralateral side (< 2cm of atrophy)
- Able to perform single leg squat with correct form
- LSI SL hop $\geq 95\%$
- Y-Balance $\geq 95\%$ (mean of 3 trials in anterior, posterolateral, posteromedial/100)
- Vertical jump test, single leg hop distance, and timed single leg hop over 20 -feet $\geq 90\%$ of contralateral side

